

Pacific Pharmaceuticals Ltd.

Lahore, Pakistan

CYSTALGON



Flavoxate HCl 200mg + Propyphenazone 250mg

Brings your Patient out of this painful and distressing state:

CYSTALGON in Urology

Effectively Manages,

- ❖ Painful Urinary Incontinence
- ❖ Hypertonia
- ❖ Incontinence due to Prostatic Obstruction / Prostatitis
- ❖ Painful Voiding, Cystitis, Urethritis etc.
- ❖ Nocturia
- ❖ Dysuria
- ❖ Painful Vesico-Urethral Spasm due to catheterization.

CYSTALGON

Suitable for both Male & Female in the treatment of painful spastic conditions
Of URO-GENITAL tract.

In infectious states; a suitable anti-infective drug should be added.

CYSTALGON in Gynaecology

Effectively Manages,

- ❖ Postpartum painful spastic conditions
- ❖ Pelvic inflammatory diseases
- ❖ Dysmenorrhoea
- ❖ Hypertonia



DOSAGE

In mild - moderate cases 1 + !

&

In moderate - severe cases 1 + 1 + 1

PRESCRIBING INFORMATION: CYSTALGON

DESCRIPTION: Cystalgon contains 200mg of Flavoxate Hydrochloride and 250mg of Propyphenazone per tablet. Cystalgon combines the properties of Flavoxate, a Recordati original specialty with marked genito-urinary anti spastic activity, and those of propyphenazone, known for its analgesic and anti-inflammatory activity.

INDICATIONS: Treatment of the following symptoms: Dysuria, Urgency, Nocturia, Suprapubic pain, incontinence in bladder and prostatic diseases such as cystitis, cystalgia, prostatitis, urethritis, urethrocystitis, inflammation of the urogenital diaphragm. Coadjuvant for analgesic-anti spastic treatment in renal and ureteral calculosis and painful urinary spastic disorders after catheterization

Pacific Pharmaceuticals Ltd.

Lahore, Pakistan

CYSTALGON



Flavoxate HCl 200mg + Propyphenazone 250mg

or cystoscopy. Aftermaths of lower urinary tract surgery. Treatment of spasm and pain in the female genital tract, for example, pelvic pain, dysrnenorrhoea, hypertonia, uterine dyskinesia.

CONTRA-INDICATIONS: Pyloric or duodenal obstructions, obstructive intestinal lesions or ileus, achalasia, gastrointestinal hemorrhage. Allergy to pyrazolonic drugs, granulocytopenia, acute intermittent porphyria, lack of glucose-6-phosphate dehydrogenase.

WARNINGS: On account of its Propyphenazone content, Cystalgon when given at high doses for prolonged period, may occasionally give rise to hypersensitivity reaction which may take the form of alterations to the skin and mucous membrane reduction in the white blood cell count, and in very rare cases - agranulocytosis. Reddish staining of the urine during treatment may be due to elimination of rubazonic acid. A harmless metabolite of the pyrazolone derivatives, and has no pathological significance. In patients hypersensitive to aminophenazone and its derivatives, high doses of Cystalgon for long periods may cause damage to the blood. Patients who drive vehicles or work with machinery, or whose normal activities call for prompt actions, should be warned of the possibility of side effects such as drowsiness, or visual disturbances.

USE IN PREGNANCY: Although tests in animals have not shown up any harmful effects on the fetus, it is advisable to use this drug with caution in pregnant women, especially during the first trimester. At normal doses, pyrazolone derivatives pass only minimally into maternal milk but caution should naturally be employed in breast-feeding women.

ADVERSE REACTIONS: Occasional patients may report nausea or vomiting (which can be limited by taking the product on a full stomach). The propyphenazone component may give rise to gastrointestinal disturbances, Hypersensitivity reactions involving the blood (Leucopenia, rarely agranulocytosis), Hypersensitivity reactions affecting the skin and mucous membranes (rash, etc) and only very rarely, anaphylactic reactions.

DOSAGE: The product should only be used by adults, children not recommended. Generally one tablet of Cystalgon 2-3 times a day, at regular intervals, depending on the severity of the pain. The SC tablets should be taken on a full stomach. Patients must not increase the doses or take the product for longer than prescribed without consulting their physician.