

Pacific Pharmaceuticals Ltd.

Lahore, Pakistan

GENURIN



Relives spastic urnary incontinence



Each Tablet contains Flavoxate HCl 100 mg / 200 mg. (Forte), an antispasmodic, selective to urogenital Tract. Specifically indicated for the treatment **OF URINARY INCONTINENCE AND Offers prompt relief of URINARY SYMPTOMS LIKE** Frequency, Urgency, Dysuria, Nocturia, Incontinence

OFTEN ASSOCIATED WITH

Cystitis, Urethritis, Prostatitis, Pelvic inflammation, Neurogenic bladder, Urge incontinence, Sequelae of surgical and endoscopic intervention in Urology.

Genurin ® Forte

- | | |
|--|--|
| <ul style="list-style-type: none">▪ Powerful relaxant of smooth muscles▪ Acts selectively on the urinary tract specially the lower pelvic region▪ Relaxes spastic muscles▪ Regulates the tone of the smooth muscles▪ Exerts musculotropic action on the Prostate tissues | <ul style="list-style-type: none">▪ Relaxes vesical spasm▪ Increases bladder capacity▪ Improves subjective symptoms rapidly▪ Has no anticholinergic side effects▪ Allows concomitant use of antibiotic (In case of infections) |
|--|--|

Genurin ® Forte offers superior profile

	Genurin ® Forte	Oxybutynin	Tolterodine
Organ specific	YES	NO	NO
Salivation	No Effect	Reduces (causes dryness of Mouth)	Reduce (causes dryness of Mouth)
Vision	No Effect	Blurred	Blurred
Effects on other Smooth muscles	No Effect	YES	YES
Anticholinergic Effects	No Effect	YES	YES

Pacific Pharmaceuticals Ltd.

Lahore, Pakistan

GENURIN



Relives spastic urinary incontinence

Indications

For the symptomatic relief of Dysuria, urgency, Nocturia, vesical suprapubic pain, frequency and incontinence as may occur in cystitis, prostatitis, urethritis. Also indicated for the relief of vesico – urethral spasms due to catheterization, prior to cystoscopy.

Dosage

Children above 11 years and above. Genurin 100 mg: 1 – 2 Tablets, 3 times a day for as long as required.

For Adults:

Genurin 200 mg (Forte): 1 – 2 Tablets 3 times a day for as long as required.

Contraindications

Genurin is contraindicated in patients who have any of the following obstructive conditions: Pyloric or duodenal obstruction, obstructive intestinal lesions or ileus, Achasis, gastrointestinal hemorrhage and obstructive uropathies of the lower urinary tract.